INFORMATIONAL LETTER NO.1907-MC-FFS

Governor

DATE: May 17, 2018

TO: Iowa Medicaid Physicians, Dentists, Advanced Registered Nurse

Practitioners, Therapeutically Certified Optometrists, Podiatrists, Pharmacies, Home Health Agencies, Rural Health Clinics, Clinics,

Lt. Governor

Skilled Nursing Facilities, Intermediate Care Facilities, Nursing Facilities-Mental ILL, Federally Qualified Health Centers (FQHC), Indian Health Service, Maternal Health Centers, Certified Nurse Midwife, Community Mental Health, Family Planning, Residential Care Facilities, ICF/ID State

Director

and Community Based ICF/ID Providers

APPLIES TO: Managed Care, Fee-for-Service

FROM: Iowa Department of Human Services (DHS), Iowa Medicaid Enterprise (IME)

RE: Iowa Medicaid Pharmacy Program Changes High Dose Opioids

EFFECTIVE: July 1, 2018

Effective July 1, 2018, prior authorization (PA) will be required for use of high-dose opioids ≥ 200 morphine milligram equivalents (MME) per day. Patients undergoing active cancer treatment or end-of-life care will not be subject to PA criteria. The MME edit will gradually be decreased over time to 90 MME per day.

PA requests should be submitted on the <u>High Dose Opioids PA form</u>¹. PA criteria for requests for opioids \geq 200 MME is as follows:

PA is required for use of high-dose opioids ≥ 200 morphine milligram equivalents (MME) per day. See <u>CDC Guideline for Prescribing Opioids for Chronic Pain</u>². Patients undergoing active cancer treatment or end-of-life care will not be subject to the criteria below. Payment will be considered when the following is met:

- 1. Requests for non-preferred opioids meet criteria for coverage (see criteria for Long-Acting Opioids and/or Short-Acting Opioids); and
- Patient has a diagnosis of severe, chronic pain with a supporting ICD-10 code. Requests for a diagnosis of fibromyalgia or migraine will not be considered; and
- 3. Patient has tried and failed at least two nonpharmacologic therapies (physical therapy; weight loss; alternative therapies such as manipulation, massage, and acupuncture; or psychological therapies such as cognitive behavior therapy [CBT]); and

¹ http://www.iowamedicaidpdl.com/pa forms

² https://www.cdc.gov/drugoverdose/prescribing/guideline.html

- 4. Patient has tried and failed at least two nonopioid pharmacologic therapies (acetaminophen, NSAIDs, or selected antidepressants and anticonvulsants; and
- 5. There is documentation demonstrating an appropriate upward titration or an appropriate conversion from other opioid medications; and
- 6. Pain was inadequately controlled at the maximum allowed dose without PA for the requested opioid(s); and
- 7. Pain was inadequately controlled by two other chemically distinct preferred long-acting opioids at the maximum allowed dose without PA; and
- 8. Chart notes from a recent office visit for pain management is included documenting the following:
 - a. Treatment plan including all therapies to be used concurrently (pharmacologic and non-pharmacologic); and
 - b. Treatment goals; and
- 9. Patient has been informed of the risks of high-dose opioid therapy; and
- 10. The prescriber has reviewed the patient's use of controlled substances on the <u>lowa</u>

 <u>Prescription Monitoring Program website</u>³ and determined that use of high-dose opioid therapy is appropriate for this patient; and
- 11. The patient's risk for opioid addiction, abuse and misuse has been reviewed and prescriber has determined the patient is a candidate for high-dose opioid therapy; and
- 12. A signed chronic opioid therapy management plan between the prescriber and patient dated within 12 months of this request is included; and
- 13. The requested dosing interval is no more frequent than the maximum FDA-approved dosing interval; and
- 14. Patient has been provided a prescription for a preferred naloxone product for the emergency treatment of an opioid overdose; and
- 15. Patient has been educated on opioid overdose prevention; and
- 16. Patient's household members have been educated on the signs of opioid overdose and how to administer naloxone; and
- 17. Patient will not be using opioids and benzodiazepines concurrently or a taper plan to discontinue the benzodiazepine must be submitted with initial and subsequent requests; and
- 18. A documented dose reduction is attempted at least annually.

If criteria for coverage are met, initial requests will be given for three months. Requests for continuation of high-dose opioid therapy will be considered every six months with the following:

- 1. High-dose opioid therapy continues to meet treatment goals, including sustained improvement in pain and function; and
- 2. Patient has not experienced an overdose or other serious adverse event; and

_

³ https://iowa.pmpaware.net/login

- 3. Patient is not exhibiting warning signs of opioid use disorder; and
- 4. The benefits of opioids continue to outweigh the risks; and
- 5. A documented dose reduction has been attempted at least annually, and the prescriber has determined the dose cannot be reduced at this time; and
- 6. The prescriber has reviewed the patient's use of controlled substances on the <u>lowa</u>

 <u>Prescription Monitoring Program website</u>⁴ and determined that continued use of highdose opioid therapy is appropriate for this patient; and
- 7. Patient will not be using opioids and benzodiazepines concurrently or a taper plan to discontinue the benzodiazepine must be submitted with subsequent requests.
- 8. Patient has been provided a prescription for a preferred naloxone product for the emergency treatment of an opioid overdose; and
- 9. Patient has been reeducated on opioid overdose prevention; and
- 10. Patient's household members have been reeducated on the signs of opioid overdose and how to administer naloxone.

We encourage providers to go to the <u>PDL website</u>⁵ to view all recent changes to the PDL. If you have questions, please contact the Pharmacy Prior Authorization Helpdesk at 877-776-1567 or 515-256-4607 (local in Des Moines) or e-mail <u>info@iowamedicaidpdl.com</u>.

⁴ https://iowa.pmpaware.net/login

⁵ http://www.iowamedicaidpdl.com/